

Journal Pre-proofs

Microwave pretreatment of sprouted mung beans before hot-air and infrared drying process

Fakhreddin Salehi

DOI: [https://doi.org/ 10.22104/IFT.2024.6635.2164](https://doi.org/10.22104/IFT.2024.6635.2164)

To appear in: *Innovative Food Technologies* (IFT)

Received Date: 7 December 2023


Revised Date: 11 January 2024

Accepted Date: 16 January 2024

Please cite this article as: Fakhreddin Salehi, Microwave pretreatment of sprouted mung beans before hot-air and infrared drying process, *Innovative Food Technologies* (2023), doi: <https://doi.org/10.22104/IFT.2024.6635.2164>

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2023 The Author(s). Published by irost.org.



Microwave pretreatment of sprouted mung beans before hot-air and infrared drying process

Fakhreddin Salehi^{1†}

¹ Associate Professor, Department of Food Science and Technology, Bu-Ali Sina University, Hamedan, Iran

Abstract

Sprouting enhances the nutritional and quality characteristics of mung beans (*Vigna radiata* L.). In this study impacts of infrared radiation and hot-air drying on microwave-pretreated mung bean sprouts were examined. The microwave process was performed for 0, 15, 30, 45, and 60 s before hot-air and infrared dehydration of sprouted mung beans. Microwave pretreatment decreased drying time of sprouted mung beans. Increasing the microwave pretreatment time from 0 to 60 s caused drying time of samples inside the hot-air dryer to decrease from 205 min to 130 min ($p < 0.05$). Drying time of sprouts in the infrared dryer was significantly less than the hot-air dryer ($p < 0.05$). Effective moisture diffusivity coefficient (D_{eff}) calculated by Fick's second law, varied in the range of 1.06×10^{-10} - 1.60×10^{-10} $m^2 s^{-1}$, and 0.98×10^{-9} - 1.15×10^{-9} $m^2 s^{-1}$, for hot-air and infrared dried samples, respectively. Experimental data for drying curves were fitted to various thin-layer equations, and the Midilli equation was best suited to explain drying kinetics of sprouted mung beans. Average rehydration ratio of dried sprouted mung beans in hot-air and infrared dryers were 262.55%, and 211.65%, respectively. In summary, the microwave pretreatment (about 60 sec) and use of infrared dryer is appropriate drying technique for sprouted mung beans with faster mass transfer and shorter drying time.

Keywords: Effective moisture diffusivity coefficient, Microwave, Midilli equation, Mung bean, Rehydration.

[†] Corresponding author email: F.Salehi@Basu.ac.ir

1. Introduction

Microwave radiation has the ability to heat a material with dielectric properties. Material absorbs microwave energy and then converts it into heat [1]. Compared to the traditional heating process, Microwave heating technology offers numerous advantages such as rapid heat transfer, non-contact heating, internal and volume heating, selective heating, uniform heating, cleanliness, pollution-free operation, and automated control [2]. Microwave drying can increase drying efficiency and reduce operating costs while maintaining high product quality [1]. Of all accessible pretreatment approaches for food products processing, use of microwaves is a promising technique [3-6]. The microwave pretreatment can facilitate mass transfer during drying and reduce drying time of fruits and vegetables [5-8]. Salehi et al. [9] used microwave pretreatment before the drying process to improve the quality attributes of dried sweet cherries. Their results showed that the moisture diffusivity, total phenolics, and antioxidant capacity of treated sweet cherries were higher than those of non-treated samples.

Mung bean (*Vigna radiata* L.) is a green legume rich in protein, oligosaccharides and fibers, high levels of phenolics, bioactive phytochemicals, flavonoids (vitexin and isovitexin), amino acids and antioxidants (regulating cholesterol levels and scavenging free radicals), and phytonutrients [10-12]. The boiled and sprouted mung beans are commonly used as a sprouted mung salad [10]. The results of Javed et al. [13] work showed that the mung bean can be utilized in production of ready-to-eat therapeutic food due to its shelf-stability and help to mitigate the protein-energy malnutrition. Liu et al. [14] and Wan et al. [15] confirmed that the mung bean or its extract can be used as an appropriate candidates of natural antioxidant and phenolic compounds.

Sprouting a traditional method used for reducing most of anti-nutritional factors in legumes. Sprouting enhances the nutritional and quality characteristics of mung beans [16, 17]. The sprouted mung beans is a good source of bioactive compounds with high antioxidant capacity [18, 19]. Dried sprouted grain powder is known to be an excellent source of nutrients for improving the quality and physicochemical properties of foods available throughout the year (such as noodles, pasta, bread, sausages, etc.) [16, 20-22]. Impacts of ultrasonic pretreatment and drying approaches on the drying kinetics and rehydration of sprouted mung beans was studied by Salehi [23]. The results demonstrated that the ultrasonic pretreatment increased the moisture diffusion capacity (higher moisture loss) and reduced the drying time of sprouted mung beans.

The use of microwaves shows many advantages. The most important aspect is shortening the time of the thermal process and reducing energy consumption and costs of the operation [1, 24]. No report on effects of microwave pretreatment on hot-air and infrared drying rate of sprouted mung beans was found. In view of the above, the aim of the study was to determine the influence of microwave pretreatment and drying approaches on drying time, mass transfer kinetic, D_{eff} , and rehydration of sprouted mung beans. In addition, the moisture ratio changes of sprouted mung beans when dried were modeled.

2. Materials and methods

2.1. Sprouting process

Mung bean seeds were purchased at a market in Arak, Markazi Province, Iran. The seeds are manually cleaned and dust-free and then stored in a dry, cool place until needed. The seeds were washed and soaked in tap water at $25\pm 1^\circ\text{C}$ for 24 hours (seeds to water ratio 1:4, w/v) [16]. Soaked seeds were kept inside a polyethylene container covered with a clean kitchen towel and allowed to germinate for 72 hours in the dark at room temperature ($25\pm 1^\circ\text{C}$). The moisture content of mung beans were determined using an oven at 105°C for 5 hours (55L, Shimaz, Iran). In this study, the average moisture content of non-treated mung beans was 7.1% (wet basis).

2.2. Microwave pretreatment

To apply the microwave pretreatment on the germinated seeds, a microwave oven (Delmonti, Model; DL740, Italy) was used under atmospheric pressure (Figure 1). In this study, the impact of the microwave pretreatment time at 5 time intervals of 0, 15, 30, 45, and 60 s on the sprouted mung beans was studied.




Figure 1. Schematic of microwave pretreatment, hot-air and infrared drying, and rehydration process of sprouted mung beans.

2.3. Hot-air drying

After each treatment, the sprouted mung beans were dried in a convective hot-air oven (at $70 \pm 2^\circ\text{C}$, 55L, Shimaz, Iran) [25].

2.4. Infrared drying

In this work, an infrared dryer (length 440 mm, width 200 mm, and height 400 mm) with an infrared radiation source (250 W, near-infrared, Noor Lamp Company, Iran) was employed for drying germinated mung beans. In this dryer, the distance from the radiant lamp to the germinated seeds was 5 cm. Mass changes of samples were recorded by a Lutron GM-300p digital balance (sensitivity of ± 0.01 g, Taiwan).

2.5. Drying kinetics

The sprouted mung beans drying kinetics are described using nine simplified drying equations (Approximation of diffusion, Henderson and Pabis, Logarithmic, Midilli, Newton, Page, Two-term, Verma, and Wang and Singh) [26, 27]. Regression analyzes were performed to estimate model parameters using Matlab software (version R2012a).

2.6. Calculation of effective moisture diffusivity coefficient (D_{eff})

The D_{eff} values (m^2/s) of sprouted mung beans during drying process, at various hot-air and infrared drying conditions, were calculated pursuant to the method explained by Salehi et al. [9].

2.7. Rehydration

Rehydration tests were performed using a water-bath (R.J42, Pars Azma Co., Iran). The dried sprouted mung beans was weighed and placed in distilled water at 50°C for 30 minutes in a 250ml beaker. I then drained the excess water for 30 seconds and reweighed the sample. The percent rehydration value of sprouted mung beans was determined as the rate of the mass of rehydrated sprouted mung beans to the dry mass of sprouted mung beans $\times 100$ [28].

2.8. Statistical analysis

The sprouted mung beans drying tests were performed with a 2 \times 5 \times 3 factorial design (dryer type, microwave treatment time, and three replicates) in a fully randomized design. Means were also compared with p -value <0.05 using Duncan's multiple range test (using the SAS 9.1, USA).

3. Results and discussion

3.1. Drying time

In this study, the final moisture content of dried sprouted mung beans was 0.30 ± 0.07 (dry basis). Enhancing air temperature in the presence of infrared radiation improved the drying rate [29]. The analysis of variance results showed that the dryer type (hot-air and infrared), microwave time, and its interactions, have a significant effects on the drying duration of sprouted mung beans ($p<0.01$) (Table 1). The impact of microwave pretreatment time and dryer type on sprouted mung beans drying duration of is presented in Figure 2. The drying time of samples dried in infrared dryer was much shorter than the samples dried in hot-air dryer at all conditions ($p<0.05$). The average drying time of sprouted mung beans dried in hot-air and infrared dryers were 171.0, and 22.0 min, respectively. The drying duration of sprouted mung beans was shortened by increasing the microwave pretreatment time. As the microwave pretreatment time increased from 0 to 60 seconds, the drying time of sprouted mung beans in the hot-air dryer decreased from 205 min to 130 min ($p<0.05$). Also, by increasing the microwave pretreatment time from 0 to 60 s, the drying time of samples inside the infrared dryer, decreased from 23.0 min to 21.0 min ($p>0.05$).

Table 1- Analysis of variance for drying time of sprouted mung beans.

Sources of changes	df	Sum of squares	Mean square	P
Dryer (D)	1	166508	166508	0.000
Microwave time (M)	4	5600	1400	0.000
D \times M	4	4830	1208	0.000
Error	20	1226	61	
Total	29	178164		




Figure 2- Effect of microwave pretreatment on drying time of sprouted mung beans (HA=Hot-air dryer; IR=Infrared dryer).

The interaction between microwaves and materials occurs through the generation of dipoles within the material and changes in their orientation. This unique feature allows simultaneous heating throughout the body, resulting in fast and uniform temperature elevation [1, 2]. The impacts of microwave time on the moisture content changes (dry basis) of sprouted mung beans when dried in the hot-air and infrared dryers are shown in Figures 3 and 4, respectively. As seen in these figures, the application of microwave has increased the rate of moisture removal from sprouted mung beans and, as a result, it increased the drying rate of the samples. Sarjerao et al. [30] used a tray dryer and a vacuum dryer to dry the sprouted mung beans. Their results confirmed that the dehydration duration in the vacuum dryer was 21.05% shorter than the drying duration in the tray dryer at 50°C.




Figure 3- Moisture content changes of sprouted mung beans when dried in the hot-air dryer




Figure 4- Moisture content changes of sprouted mung beans when dried in the infrared dryer

3.2. Effective moisture diffusivity coefficient (D_{eff})

In constant power microwave drying, as the moisture content in the material decreases, its ability to absorb microwaves decreases [24]. The impacts of microwave pretreatment time and dryer type on the D_{eff} values of sprouted mung beans are presented in Figure 5. The D_{eff} values of the samples dried with the infrared dryer were considerably higher than those of the sprouts dried with the hot-air dryer ($p < 0.05$). The average D_{eff} values of sprouted mung beans when dried in hot-air dryer increased from $1.06 \times 10^{-10} \text{ m}^2 \text{ s}^{-1}$ to $1.60 \times 10^{-10} \text{ m}^2 \text{ s}^{-1}$ when the microwave pretreatment time was increased from 0 to 60 s ($p > 0.05$). In addition, the average D_{eff} values of sprouted mung beans when dried in infrared dryer increased from $0.98 \times 10^{-9} \text{ m}^2 \text{ s}^{-1}$ to $1.15 \times 10^{-9} \text{ m}^2 \text{ s}^{-1}$ when the microwave pretreatment time was increased from 0 to 60 s ($p < 0.05$). These data lie within the common range of 10^{-11} – $10^{-9} \text{ m}^2/\text{s}$ for other grains. Rafiee et al. [31] used a hot-air dryer (at 35–70°C) for drying wheat (Tajan). The results indicated that the D_{eff} values for the whole wheat ranged from $2.3 \times 10^{-11} \text{ m}^2/\text{s}$ to $1.1 \times 10^{-10} \text{ m}^2/\text{s}$, and it was enhanced when the temperature of the dry air increased. Results of Sarjerao et al. [30] study indicated that the D_{eff} values for vacuum-dried sprouted mung beans was more than tray dryer at all condition. Also, the D_{eff} values for tray and vacuum dryers were reported between $3.98 \times 10^{-11} \text{ m}^2 \text{ s}^{-1}$ and $7.35 \times 10^{-11} \text{ m}^2 \text{ s}^{-1}$, and $4.50 \times 10^{-11} \text{ m}^2 \text{ s}^{-1}$ and $7.892 \times 10^{-11} \text{ m}^2 \text{ s}^{-1}$, respectively. Manikantan et al. [27] used a tray dryer (at 50–80°C, for 24–48 h) for drying sprouted wheat grains. Their results indicated that the D_{eff} values of sprouted wheat grains (at 48h drying period) increased from $1.9 \times 10^{-9} \text{ m}^2 \text{ s}^{-1}$ to $2.6 \times 10^{-9} \text{ m}^2 \text{ s}^{-1}$ with an increase in dryer temperature from 50 to 80°C. In another study, Shingare and Thorat [22] used a fluidized bed dryer to dry sprouted wheat grains. They reported D_{eff} values for sprouted wheat grains when dried in a fluidized bed dryer ranging from $7.3 \times 10^{-10} \text{ m}^2 \text{ s}^{-1}$ to $30.4 \times 10^{-10} \text{ m}^2 \text{ s}^{-1}$.




Figure 5- Effect of microwave pretreatment on effective water diffusivity coefficient of sprouted mung beans (HA=Hot-air dryer; IR=Infrared dryer).

3.3. Kinetics modelling

The drying behavior of sprouted mung beans in hot-air and infrared dryers was fitted with the Midilli model. Combining the highest r value with the lowest sum of squares for error (SSE) and root mean squared error (RMSE) values establishes a satisfactory agreement between the actual data and the correlations. This model presented a good fit with the maximum r -value (greater than 0.998) and the

minimum SSE, and RMSE values (less than 0.0053 and 0.0219, respectively) for all conditions. The calculated constant coefficients of the Midilli equation include a, k, n, and b, are shown in Table 2. The values of SSE, RMSE, and r values for all conditions ranged from 0.0004 to 0.0053, 0.0036 to 0.0219, and 0.998 to 0.999, respectively. Figure 6 shows a comparison of the experimental results (microwave time = 30 s, infrared dryer used) and fitted moisture ratio data using the Midilli equation. The results demonstrate that the Midilli equation is suitable for describing the drying rate of sprouted mung beans under different drying conditions. The results of Salehi [23] study indicated that Midilli was the appropriately fitted equation for sprouted mung beans drying in the hot-air and infrared dryers.

Table 2- The constants and coefficients of the Midilli model

Dryer type	Microwave time (sec)	a	k	n	b	Sum of squared error (SSE)	Correlation coefficient (r)	Root mean square error (RMSE)
Hot-air dryer	0	0.9951	0.0060	0.8803	-0.0045	0.0022	0.9990	0.0141
	15	0.9887	-0.0005	1.2211	-0.0039	0.0015	0.9993	0.0114
	30	0.9864	0.0043	1.2280	0.0000	0.0027	0.9989	0.0157
	45	0.9818	0.0012	1.5397	0.0005	0.0025	0.9990	0.0147
	60	0.9815	0.0019	1.4573	0.0002	0.0042	0.9985	0.0194
Infrared dryer	0	1.0103	-0.0199	1.2566	-0.0403	0.0017	0.9995	0.0075
	15	1.0026	0.0301	1.3617	0.0010	0.0005	0.9998	0.0041
	30	1.0047	0.0186	1.2197	-0.0273	0.0007	0.9996	0.0046
	45	1.0147	0.0357	1.3973	0.0016	0.0008	0.9995	0.0052
	60	1.0021	0.0372	1.2703	-0.0005	0.0012	0.9997	0.0063




Figure 6- Evaluation of fitted values by Midilli model with experimental data of moisture ratio (microwave pretreatment time=30 s and at infrared dryer).

3.4. Rehydration

Analysis of variance results confirmed that the drying method (hot-air and infrared), microwave time, and their interaction had a significant influence on the rehydration rate of dried sprouted mung beans ($p < 0.01$) (Table 3). The impacts of microwave pretreatment and drying approaches on the rehydration ratio of dried sprouted mung beans are shown in Figure 7. The rehydration rate of dried sprouted mung beans in hot-air and infrared dryers increased when the microwave pretreatment time was increased. Furthermore, the rehydration rate of the pretreated hot-air-dried sprouted mung beans was considerably higher than that of the infrared dried sprouted mung beans ($p < 0.05$). This problem may be due to the small shrinkage, large volume and porous structure of the hot-air dried sprouts, resulting in greater water diffusion within the dried mung bean cells and a more rehydration rate [23]. The mean rehydration rate of dried sprouted mung beans in the hot-air dryer enhanced from 254.85% to 273.72% as the microwave pretreatment time was increased from 0 to 60 seconds ($p < 0.05$). In addition, the average rehydration ratio of dried sprouted mung beans in infrared dryer increased from 187.60% to 222.48% when the microwave pretreatment time was increased from 0 to 60 seconds ($p < 0.05$). Salehi [23] reported that the rehydration ratio of dried sprouted mung beans in hot-air and infrared dryers increased when the ultrasound pretreatment time was increased.

Table 3- Analysis of variance for rehydration of dried sprouted mung beans in hot-air and infrared dryers.

Sources of changes	df	Sum of squares	Mean square	P
Dryer (D)	1	19431.5	19431.5	0.000
Microwave time (M)	4	2531.8	633.0	0.000
D × M	4	621.1	155.3	0.000
Error	20	305.0	15.3	
Total	29	22889.4		




Figure 7- Effect of microwave pretreatment on rehydration of dried sprouted mung beans (HA=Hot-air dryer; IR=Infrared dryer).

4. Conclusion

In this work, I examined the impacts of microwave pretreatment and drying approach on the thin layer dehydration properties of sprouted mung beans. The drying time of samples dried with an infrared dryer was considerably shorter than that of sprouts dried with a hot-air dryer ($p < 0.05$). The average D_{eff} values of sprouted mung beans when dried in the hot-air and infrared dryers enhanced when the microwave pretreatment duration was increased from 0 to 60 s. The average D_{eff} values of sprouted mung beans when dried in hot-air and infrared dryers were $1.28 \times 10^{-10} \text{ m}^2 \text{ s}^{-1}$, and $1.04 \times 10^{-9} \text{ m}^2 \text{ s}^{-1}$, respectively. The drying behavior of sprouted mung beans was fitted using the Midilli equation. This formula showed a good fit with the maximum r-value (greater than 0.998), and the minimum SSE and RMSE values (less than 0.0053 and 0.0219, respectively) for all conditions compared to other models. The rehydration rate of pretreated and hot-air dried sprouts was considerably higher than that of infrared dried sprouts ($p < 0.05$). The results of this work confirmed that microwave pretreatment improved the mass transfer rate of sprouted mung beans when dried in hot-air and infrared dryers.

References

- [1] Guzik, P., Kulawik, P., Zając, M., & Migdał, W. (2022). Microwave applications in the food industry: an overview of recent developments. *Crit. Rev. Food Sci. Nutr.*, 62, 7989-8008. doi: 10.1080/10408398.2021.1922871
- [2] Ren, C., Zhou, J., Zhang, Y., Omran, M., Gao, L., Tang, J., Zhang, F., & Chen, G. (2024). Thin layer modeling the drying kinetics of titanium slag under microwave-assisted drying. *Adv. Powder Technol.*, 35, 104312. doi: 10.1016/j.apt.2023.104312
- [3] Singh, S., Bhat, H.F., Kumar, S., Lone, A.B., Aadil, R.M., Ait-Kaddour, A., Hassoun, A., Proestos, C., & Bhat, Z.F. (2023). Ultrasonication and microwave pre-treated locust protein hydrolysates enhanced the storage stability of meat emulsion. *Ultrason. Sonochem.*, 98, 106482. doi: 10.1016/j.ultsonch.2023.106482
- [4] Xu, J. (2015) Chapter 9-Microwave pretreatment, in: A. Pandey, S. Negi, P. Binod, C. Larroche (Eds.) *Pretreatment of Biomass*, Elsevier, Amsterdam, pp. 157-172.
- [5] Mothibe, K.J., Zhang, M., Mujumdar, A.S., Wang, Y.C., & Cheng, X. (2014). Effects of ultrasound and microwave pretreatments of apple before spouted bed drying on rate of dehydration and physical properties. *Drying Technol.*, 32, 1848-1856. doi: 10.1080/07373937.2014.952381
- [6] Wray, D., & Ramaswamy, H.S. (2015). Novel concepts in microwave drying of foods. *Drying Technol.*, 33, 769-783. doi: 10.1080/07373937.2014.985793
- [7] Sharma, G.P., & Prasad, S. (2006). Optimization of process parameters for microwave drying of garlic cloves. *J. Food Eng.*, 75, 441-446. doi: 10.1016/j.jfoodeng.2005.04.029
- [8] Salehi, F., Goharpour, K., & Razavi Kamran, H. (2024). Effects of different pretreatment techniques on the color indexes, drying characteristics and rehydration ratio of eggplant slices. *Results Eng.*, 21, 101690. doi: 10.1016/j.rineng.2023.101690
- [9] Salehi, F., Inanloodoghuz, M., & Ghazvineh, S. (2023). Influence of microwave pretreatment on the total phenolics, antioxidant activity, moisture diffusivity, and rehydration rate of dried sweet cherry. *Food Sci. Nutr.*, 11, 7870-7876. doi: 10.1002/FSN3.3703
- [10] Liyanage, R., Kiramage, C., Visvanathan, R., Jayathilake, C., Weththasinghe, P., Bangamuwage, R., Chaminda Jayawardana, B., & Vidanarachchi, J. (2018). Hypolipidemic and hypoglycemic potential of raw, boiled, and sprouted mung beans (*Vigna radiata* L. Wilczek) in rats. *J. Food Biochem.*, 42, e12457. doi: 10.1111/jfbc.12457
- [11] Bai, Y., Chang, J., Xu, Y., Cheng, D., Liu, H., Zhao, Y., & Yu, Z. (2016). Antioxidant and myocardial preservation activities of natural phytochemicals from mung bean (*Vigna radiata* L.) seeds. *J. Agric. Food Chem.*, 64, 4648-4655. doi: 10.1021/acs.jafc.6b01538
- [12] Tang, D., Dong, Y., Ren, H., Li, L., & He, C. (2014). A review of phytochemistry, metabolite changes, and medicinal uses of the common food mung bean and its sprouts (*Vigna radiata*). *Chem. Cent. J.*, 8, 1-9. doi: 10.1186/1752-153X-8-4
- [13] Javed, F., Jabeen, S., Sharif, M.K., Pasha, I., Riaz, A., Manzoor, M.F., Sahar, A., Karrar, E., & Aadil, R.M. (2021). Development and storage stability of chickpea, mung bean, and peanut-based ready-to-use therapeutic food to tackle protein-energy malnutrition. *Food Sci. Nutr.*, 9, 5131-5138. doi: 10.1002/fsn3.2479
- [14] Liu, D., Guan, X., Huang, K., Li, S., Liu, J., Yu, W., & Duan, R. (2019). Protective effects of mung bean (*Vigna radiata* L.) and pea (*Pisum sativum* L.) against high-fat-induced oxidative stress. *Food Sci. Nutr.*, 7, 4063-4075. doi: 10.1002/fsn3.1271

- [15] Wan, F., Hou, C., Luo, K., & Cheng, A. (2022). Steam explosion enhances phenolic profiles and antioxidant activity in mung beans. *Food Sci. Nutr.*, 10, 1039-1050. doi: 10.1002/fsn3.2711
- [16] El-Adawy, T.A., Rahma, E.H., El-Bedawey, A.A., & El-Beltagy, A.E. (2003). Nutritional potential and functional properties of germinated mung bean, pea and lentil seeds. *Plant Foods Hum. Nutr.*, 58, 1-13. doi: 10.1023/B:QUAL.0000040339.48521.75
- [17] Sangsukiam, T., & Duangmal, K. (2017). A comparative study of physico-chemical properties and antioxidant activity of freeze-dried mung bean (*Vigna radiata*) and adzuki bean (*Vigna angularis*) sprout hydrolysate powders. *Int. J. Food Sci. Tech.*, 52, 1971-1982. doi: 10.1111/ijfs.13469
- [18] Wei, Y., Wang, X., Shao, X., Xu, F., & Wang, H. (2019). Sucrose treatment of mung bean seeds results in increased vitamin C, total phenolics, and antioxidant activity in mung bean sprouts. *Food Sci. Nutr.*, 7, 4037-4044. doi: 10.1002/fsn3.1269
- [19] Elobuiké, C.S., Idowu, M.A., Adeola, A.A., & Bakare, H.A. (2021). Nutritional and functional attributes of mungbean (*Vigna radiata* [L] Wilczek) flour as affected by sprouting time. *Legume Science*, 3, e100. doi: 10.1002/leg3.100
- [20] Jokar, A., NoruziPaghband, A., Madani, S., Shaamirian, M., & Zare, M. (2019). Using germinated wheat flour instead of flour and starch in sausage production. *J. Food Sci. Technol. (Iran)*, 15, 61-72.
- [21] Liu, Y., Xu, M., Wu, H., Jing, L., Gong, B., Gou, M., Zhao, K., & Li, W. (2018). The compositional, physicochemical and functional properties of germinated mung bean flour and its addition on quality of wheat flour noodle. *J. Food Sci. Technol.*, 55, 5142-5152. doi: 10.1007/s13197-018-3460-z
- [22] Shingare, S.P., & Thorat, B.N. (2013). Fluidized bed drying of sprouted wheat (*Triticum aestivum*). *Int J Food Eng*, 10, 29-37. doi: 10.1515/ijfe-2012-0097
- [23] Salehi, F. (2023). Effects of ultrasonic pretreatment and drying approaches on the drying kinetics and rehydration of sprouted mung beans. *Legume Science*, 5, e211. doi: 10.1002/leg3.211
- [24] Wu, X., Zhao, W., Wang, X., Bai, Z., & Ma, L. (2023). A novel variable power microwave (VPM) drying technology for lowering energy consumption and improving the in vitro protein digestibility of black soldier fly larvae. *Innov. Food Sci. Emerg. Technol.*, 89, 103470. doi: 10.1016/j.ifset.2023.103470
- [25] Salehi, F., & Inanloodoghous, M. (2023). Effects of gum-based coatings combined with ultrasonic pretreatment before drying on quality of sour cherries. *Ultrason. Sonochem.*, 100, 106633. doi: 10.1016/j.ultsonch.2023.106633
- [26] Salehi, F., & Satorabi, M. (2021). Influence of infrared drying on drying kinetics of apple slices coated with basil seed and xanthan gums. *Int. J. Fruit Sci.*, 21, 519-527. doi: 10.1080/15538362.2021.1908202
- [27] Manikantan, M., Mridula, D., Sharma, M., Kochhar, A., Prasath, V.A., Patra, A., & Pandiselvam, R. (2022). Investigation on thin-layer drying kinetics of sprouted wheat in a tray dryer. *Qual. Assur. Saf. Crop. Foods*, 14, 12-24.
- [28] Salehi, F., Razavi Kamran, H., & Goharpour, K. (2023). Effects of ultrasound time, xanthan gum, and sucrose levels on the osmosis dehydration and appearance characteristics of grapefruit slices: process optimization using response surface methodology. *Ultrason. Sonochem.*, 98, 106505. doi: 10.1016/j.ultsonch.2023.106505
- [29] Salehi, F. (2020). Recent applications and potential of infrared dryer systems for drying various agricultural products: A review. *Int. J. Fruit Sci.*, 20, 586-602. doi: 10.1080/15538362.2019.1616243
- [30] Sarjerao, L.S.K., Kashyap, P., & Sharma, P. (2022). Effect of drying techniques on drying kinetics, antioxidant capacity, structural, and thermal characteristics of germinated mung beans (*Vigna radiata*). *J. Food Process Eng.*, 45, e14155. doi: 10.1111/jfpe.14155
- [31] Rafiee, S., Keyhani, A., & Jafari, A. (2008). Modeling effective moisture diffusivity of wheat (Tajan) during air drying. *Int. J. Food Prop.*, 11, 223-232. doi: 10.1080/10942910701291858

Journal Pre-proofs